Neurophysiological encoding of VOT and F0 in voicing perception

Chao Han¹, Katie Ma¹, Yoonjung Kang^{1,2}, Philip J. Monahan^{1,2,3}

- Department of Language Studies, University of Toronto Scarborough
- Department of Linguistics, University of Toronto
- Department of Psychology, University of Toronto Scarborough









PREVIEW

- Both VOT and F0 support English consonant voicing categorization.
- Individuals differ in their VOT and F0 cue weights.
- MMN does not track individual VOT and F0 cue weights.

BACKGROUND

English stop voicing contrasts primarily cued by voice onset time (VOT), secondarily by fundamental frequency (F0) [1].







- Individual variation in cue weighting [2].
- Larger mismatch negativity (MMN) response linked to more robust phonological contrast encoding [3].

Research Questions:

- Do secondary cues pre-attentively engage phonological encoding?
- 2. Do MMN responses reflect participant-level variation in cue-weighting?

METHODS






RESULTS: Behavioral cue weights

Mixed-effects logistic regression for each group.

- Fixed effect: normalized cue value
- Random effects: Random intercepts and random by-participant slopes for normalized cue value

Regression coefficients extracted for each participants to index individual cue weights.

- VOT group: $\beta = 4.3$, p < .001
- F0 group: β = 2.6, p < .001
- A significant difference between VOT and FO slopes: t = 6.3, p < .001




RESULTS: MMN

- Identity MMN: subtract ERPs to same stimulus, standard presentation minus deviant presentation across blocks.
- MMN amplitude: assessed by spatiotemporal cluster-based permutation tests (cluster formation threshold p = 0.05, twotailed; cluster-level threshold: p = 0.05).

(* = significant cluster found; n.s. = no significant cluster)

	F0 group	VOT group
Dorsal	lowFO (n.s.)	shortVOT (*)
/ka/ vs. /ga/	highF0 (*)	longVOT (n.s.)
Glottal	lowF0 (n.s.)	shortVOT (n.s.)
/ha/	highFO (*)	longVOT (n.s.)



Example GAM curve


Half-are a latency

RESULTS: Brain-behavior relations

Individual brain responses for each deviant condition, modeled using generalized additive modeling (GAM) [4]. Two measures extracted:

- Normalized modeled peak: MMN peak amplitude normalized by standard error: peak / (1.96 * SE)
- Half-area latency: Time point at which 50% of the area within the negative deflection is reached.

Correlate individual responses with individual cue weights.



DISCUSSION & REFERENCES

- Behaviorally, F0 supports English voicing contrasts.
- VOT MMN found for dorsal shortVOT but not for longVOT
- Consistent with phonological accounts [5–7].
- FO MMN comparable for dorsal and glottal conditions:
- No evidence that F0 is engaged pre-attentively in phonological contrast encoding.
- No significant brain-behavior correlations.
 - Neural response encoding individual cue weights may degrade by MMN (150–300 ms).
 - Consistent with degradation of fine-phonetic details along the subcortical-cortical auditory pathway [8-10].

1. Kapnoula, E. C., Winn, M. B., Kong, E. J., Edwards, J. & McMurray, B. Evaluating the sources and functions of gradiency in phoneme categorization: An individual differences approach. J. Exp. Psychol. Hum. Percept. Perform. 43, 1594–1611 (2017). 2. Kong, E. J. & Edwards, J. Individual differences in categorical perception of speech: Cue weighting and executive function. J. Phon. 59,

3. Díaz, B., Baus, C., Escera, C., Costa, A. & Sebastián-Gallés, N. Brain potentials to native phoneme discrimination reveal the origin of individual differences in learning the sounds of a second language. Proc. Natl. Acad. Sci. 105, 16083-16088 (2008).

4. Meulman, N., Sprenger, S. A., Schmid, M. S. & Wieling, M. GAM-based individual difference measures for L2 ERP studies. Res. 5. Lahiri, A. & Reetz, H. Underspecified recognition. in Laboratory Phonology 7 (eds Gussenhoven, C. & Warner, N.) 637–676 (Mouton de Gruyter, 2002). doi:10.1515/9783110197105.2.637.

5. Hestvik, A. & Durvasula, K. Neurobiological evidence for voicing underspecification in English. *Brain Lang.* 152, 28–43 (2016). 7. Fu, Z. & Monahan, P. J. Extracting Phonetic Features From Natural Classes: A Mismatch Negativity Study of Mandarin Chinese Retroflex Consonants. Front. Hum. Neurosci. 15, 1–15 (2021). 8. Tamura, S. & Sung, Y. Brainstem and early cortical auditory activities associated with language differences in acoustic cue weighting

9. Toscano, J. C., McMurray, B., Dennhardt, J. & Luck, S. J. Continuous perception and graded categorization: Electrophysiological evidence for a linear relationship between the acoustic signal and perceptual encoding of speech. Psychol. Sci. 21, 1532–1540 (2010). 10. Ou, J., Xiang, M. & Yu, A. C. L. Individual variability in subcortical neural encoding shapes phonetic cue weighting. Sci. Rep. 13, 9991